

What is a syndrome?



A syndrome is a group of signs or symptoms that happen together and help to identify a unique medical condition. Epilepsy syndromes are usually diagnosed in childhood but some can continue into adult life.

What is an 'epilepsy syndrome'?

If your child is diagnosed with an epilepsy syndrome, it means that their epilepsy has some specific signs and symptoms. These include:

- the type of seizure or seizures they have;
- the age when the seizures start;
- a specific pattern on an electroencephalogram (EEG); and
- sometimes a pattern on a brain imaging scan.

An EEG test is painless and records patterns of electrical activity in the brain. Some epilepsy syndromes have a particular pattern, so the EEG can be helpful in finding the correct diagnosis. A magnetic resonance imaging (MRI) brain scan is also painless and looks at the structure of the brain for any underlying abnormality. Children may need a general anaesthetic to ensure they remain still during the MRI scan.

An epilepsy syndrome can only be diagnosed by looking at all the signs and symptoms together.

[Visit epilepsysociety.org.uk/diagnosis](https://epilepsysociety.org.uk/diagnosis)
[and epilepsysociety.org.uk/epileptic-seizures](https://epilepsysociety.org.uk/epileptic-seizures)

If your child is diagnosed with an epilepsy syndrome, it may help the paediatrician (a doctor who specialises in treating children) to plan their care (for example, choosing treatment options or deciding whether further tests are needed).

Different types of syndrome

Syndromes can vary greatly. Some are called 'self-limited', which means children become seizure free (have no seizures) once they reach a certain age. Other syndromes are 'severe' and children have seizures which are difficult to control. Anti-seizure medications (ASMs) may be tried alone, or in combination with each other, and some non-drug treatments may also be tried, for example the ketogenic diet.

Many children with severe epilepsy syndromes have additional difficulties with learning and behaviour and may need extra support.

[Visit epilepsysociety.org.uk/treatment](https://epilepsysociety.org.uk/treatment)

Examples of childhood syndromes

Self-limited epilepsy with centrotemporal spikes (SeLECTS)

SeLECTS, previously called benign rolandic epilepsy, affects around 15% of children with epilepsy and can start any time between the ages of 3 and 13.

Children may have very few seizures and most become seizure-free by the age of 16. They may have focal preserved consciousness seizures with observable manifestations (previously called focal motor aware seizures), which means they involve movement. They are often at night, and usually involve one side of the face and/or the muscles that involve speech and swallowing, causing gurgling or grunting noises, mouth movements, and dribbling. Speech can be temporarily affected and symptoms may develop into a tonic clonic seizure. ASMs may not be necessary, but can be helpful if seizures are more frequent or are mostly tonic clonic.

There are many different types of epilepsy syndrome. This factsheet gives a brief overview of what epilepsy syndromes are and includes details of some specific syndromes and their characteristics.

Helpline 0300 102 0024
Confidential, information, and
emotional support.
[Visit epilepsysociety.org.uk/helpline](https://epilepsysociety.org.uk/helpline)
for opening hours.

Childhood absence epilepsy (CAE)

This syndrome usually starts between the ages of four and ten, and can affect up to 18% of school age children with epilepsy. Absence seizures happen frequently (up to 100 times a day) and are very brief, lasting only a few seconds. Because of this, they often go unnoticed. During a seizure a child will become unconscious. They may look blank or stare, their eyelids may flutter, and they may make repetitive movements. They may not respond to what is happening around them, or be aware of what they are doing.

Seizures respond well to anti-seizure medication. If a child is seizure-free for two years, medication is sometimes gradually reduced. Up to 90% of children with CAE will grow out of seizures by the time they are adults. Sometimes a child with CAE may also have other types of seizures.

Dravet syndrome

Dravet syndrome is a rare and complex neurological condition that affects around one in every 15,000 babies born in the UK. Seizures usually begin in the first year of life, with additional characteristics emerging typically from the second year onwards.

The seizures can happen unexpectedly and are often linked to having a high temperature or being somewhere hot. This syndrome is often associated with difficult to treat seizures, intellectual and behavioural difficulties, and a range of other problems. Dravet syndrome is a lifelong condition, but its symptoms and effects can change over time.

[Visit epilepsysociety.org.uk/news/epilepsy-society-research-published-brain-communications-journal-and-epilepsysociety.org.uk/news/treatment-ultra-rare-form-epilepsy-shows-promising-results](https://epilepsysociety.org.uk/news/epilepsy-society-research-published-brain-communications-journal-and-epilepsysociety.org.uk/news/treatment-ultra-rare-form-epilepsy-shows-promising-results)

Juvenile myoclonic epilepsy (JME)

This syndrome usually starts between the ages of 12 and 18. Many children have different types of seizure: myoclonic seizures (brief muscle jerks) in the upper body and generalised tonic clonic seizures. Some children may also have occasional absence seizures.

Seizures often happen as, or shortly after waking. Medication can be successful in controlling seizures, and may be needed for life. Tiredness, stress, lack of sleep, and excess alcohol can trigger seizures. It is common for people with JME to have photosensitive epilepsy.

[Visit epilepsysociety.org.uk/photosensitive-epilepsy](https://epilepsysociety.org.uk/photosensitive-epilepsy)

Infantile spasms (or West Syndrome)

This syndrome often begins in the first year of life and can affect children:

- who have had a previous brain injury before the age of six months;
- whose brain has not formed properly (brain malformation); or
- who have genetic abnormalities.

It is identified by brief spasms or jerks which happen in 'clusters'. Spasms can affect the whole body or just the arms and legs. Each cluster can include between 10 – 100 individual spasms, which often happen when the child is waking up. ASMs and corticosteroids (medicines to reduce inflammation) are used to treat this syndrome, and medicines help to control spasms in about seven out of ten children. Many children develop problems with learning or behaviour. Some may go on to develop Lennox-Gastaut syndrome.

Lennox-Gastaut syndrome

This syndrome usually begins between the ages of three and five, but can start as late as adolescence. Children may have different types of seizures, most commonly tonic (where muscles become stiff), atonic (where muscles relax), and atypical absences.

Atypical absences are different from typical absences as they often last longer, and a child may be responsive and aware of their surroundings.

Many children also develop learning difficulties and behaviour problems. This syndrome can be very difficult to treat with ASM, and most children need a combination of different drugs. Some non-drug treatments, such as the ketogenic diet and vagus nerve stimulation therapy (VNS), can also be helpful. Seizures often continue into adult life.

[Visit epilepsysociety.org.uk/treatment-and-epilepsysociety.org.uk/news/life-changing-treatment-lennox-gastaut-syndrome](https://epilepsysociety.org.uk/treatment-and-epilepsysociety.org.uk/news/life-changing-treatment-lennox-gastaut-syndrome)

Who can I talk to?

If your child has been diagnosed with an epilepsy syndrome, you may have concerns or questions. You can get information and advice from a paediatrician with an interest in epilepsy, or a paediatric neurologist, or from a neurologist for adults. Support may also be available through an epilepsy specialist nurse, counsellor, support group, friends, family, or a helpline (see page 1).

Every effort is made to ensure that all information is correct at the time of publishing. Please note that information is intended for a UK audience. This information is not a substitute for advice from your own doctors. Epilepsy Society is not responsible for any actions taken as a result of using this information.



**epilepsy
society**

Further information and support

Angelman Syndrome

angelmanuk.org

Angelman Syndrome support education and research trust (ASSERT).

Cambridge Rare Disease Network UK

camraredisease.org

Unites patients, advocates, experts and leaders to address the challenges faced by people affected by rare diseases.

Contact a family

contact.org.uk

Information and links to support for many childhood conditions, including epilepsy syndromes.

Corpal

corpal.org.uk

Supporting those with disorders of the corpus callosum (DCC) or Aicardi Syndrome.

Dravet Syndrome UK

dravet.org.uk

Support education and research for those affected.

USA Epilepsy Foundation rare epilepsy network

epilepsy.com/what-is-epilepsy/rare-epilepsies/ren

Epilepsy Research Institute UK

epilepsy-institute.org.uk

Advocates and actions the research priorities of people affected by epilepsy.

European Epicare

epi-care.eu

The European Reference Network for all rare and complex epilepsies.

Epilepsy Sparks

epilepsysparks.com/rare-epilepsy-organisations

List of international information/support groups.

Lennox Gastaut Syndrome

lgsfoundation.org

Support for Lennox Gastaut syndrome in the UK.

Rare epilepsy network

rareepilepsynetwork.org

Aims to improve outcomes for people with rare epilepsies, based in the USA.

Rett Syndrome

rettsyndrome.org.uk

Information on Rett Syndrome.

Ring Chromosome 20 Syndrome

ring20researchsupport.co.uk

Research, information and support for Ring Chromosome 20 Syndrome.

Sturge Weber Foundation

sturgeweber.org.uk

West Syndrome

wssg.org.uk

UK rare epilepsies together

ukret.co.uk

A UK rare epilepsies group with a network of patient advocacy groups.

Young Epilepsy

youngepilepsy.org.uk

Information and support with epilepsy syndromes.

Epilepsy Society is grateful to Dr F J Rugg-Gunn, Consultant Neurologist & Honorary Associate Professor, Clinical Lead, Chalfont Centre for Epilepsy, who reviewed this information.

For a printed copy of this information contact our helpline.

Epilepsy Society

Chesham Lane,
Chalfont St Peter,
Buckinghamshire
SL9 0RJ



@epilepsysociety



Patient Information Forum

Every effort is made to ensure that all information is correct at the time of printing. Please note that information is intended for a UK audience. This information is not a substitute for advice from your own doctors. Epilepsy Society is not responsible for any actions taken as a result of using this information.